

## **TERIPARATIDE Do(s) AND Don't(s)**

<u><b>Sr.No.</b></u>	<u><b>Do(s)</b></u>	<u><b>Don't(s)</b></u>
1.	Use the injection as prescribed by qualified dr. only	Do not use injection without prescription of qualified dr.
2.	Check the injection when delivered at your doorstep. It must be delivered in Ice pack only	Do not use injection, which has been delivered without ice pack.
3.	Use the cartridge of terifrac must be sealed and with colourless liquid inside.	Don't use the cartridge, if broken or having residue inside
4.	Dial the dosage of pen upto mark 8 (which is equivalent to 20 mcg dosage) and then inject by needle.	Don't dial more than 8 no. marked in pen, if by mistake it is more than 8 than don't reverse the no.
5.	If by mistake dose is dialed more than 8 no. in pen then dial clockwise and complete numbers and then again dial upto 8	Don't do reverse dialing if by mistake it is more than 8.
6.	Use the needle of Becton Dickinson or standard company needles from 29 to 31 gauges only	Don't use the needle without company brand or less than 29 gauges or more than 31 gauges.
7.	Store the pen between 2 to 8 degree temperature	Don't store the pen on room temperature or donot freeze it.
8.	Inject Terifrac right away after you take the delivery device out of the refrigerator.	Don't delay for injection after taking the device out of refrigerator
9.	Terifrac can be taken at any time with or without food. Take injection at the same time each day if possible.	Do not take more than one injection in the same day.
10.	If a dose is missed on a particular day have administration of injection at the next scheduled day on the same timing.	Don't have overlap of two inj. pricks with a very short interval in-between.
11.	Terifrac can be used in Postmenopausal Osteoporosis as well as Glucocortoid Induced and Hypogonadal osteoporosis as per US FDA approval	Terifrac cannot be used in the patient with Paget's disease, other bone disease, cancer in bones ,kidney stones, radiation therapy, hyperparathyroidism, women who is in breast feeding or pregnant.
12.	Terifrac can be used for 1.5 yrs to 2 yrs as per FDA guideline.	It cannot be used more than 2 years.
13.	Terifrac injection can be injected subcutaneously at abdomen or thigh part.	Don't use or inject terifrac in other parts of body except abdomen and thigh
14.	P1NP bone marker test can be done prior to taking terifrac injection, which is usually followed after 3 or 6 months.	Do not do bone marker test after starting terifrac treatment.
15.	Keep terifrac in Ice pack while travelling.	Don't keep it out of icepack for a prolonged period while travelling.